

Protocol N: for (Children) and Adolescents (Ninos is children in Spanish)

Protocol N, for children, is based on the experiences and testimonials of many mothers.

- As a general rule for an initial **hourly dose**, we use a maximum of one drop of activated CD for every 12 kilograms of body weight, in 100 ml or more of water. CDS is often tolerated better than CD; **use 1 ml of CDS 0.3% (3000 ppm) in 100 ml** or more of water for every 12 kilograms of weight.
- Before starting any protocol, first check compatibility, to avoid any adverse effects.
- Always activate the drops on a 1:1 ratio, adding between 100 and 200 ml of water.
- Watch for any adverse reactions from the minor: fatigue, nausea, belly pain, vomiting, etc. to adjust the dose accordingly.
- Except for extremely critical and urgent cases, it s best not to administer any prescription drug or treatment to children under the age of one year.

Dosage (drops according to the child's weight)

5 kilos:	3 drops a day divided into ten intakes. The drops may be mixed with rice milk.
15 kilos:	6 drops a day divided into ten intakes.
30 kilos:	8 drops a day divided into ten intakes.
40 kilos:	12 drops a day divided into ten intakes.
60 kilos:	adult dose

Notes:

- » Avoid antioxidants and Vitamin C.
- » Keep all products out of the reach of children.
- » Try to use child-proof containers.

